

Many-body theory of degenerate systems: A simple example

Christian Brouder

*Laboratoire de Minéralogie Cristallographie, CNRS UMR 7590, Universités Paris 6 et 7,
IPGP, case 115, 4 place Jussieu, 75252 Paris cedex 05, France.*

(Dated: February 2, 2008)

The hierarchy of Green functions for (quasi)degenerate systems, presented in cond-mat/0308058, is calculated in detail for the case of a system with closed shells plus a single electron in a two-fold degenerate level. The complete hierarchy is derived explicitly, in terms of Green functions and of Feynman diagrams.

I. INTRODUCTION

In reference¹, hereafter referred to as I, a many-body theory of degenerate or quasidegenerate systems was presented, using quantum group methods. In particular, a hierarchy of Green function was derived and formulated in terms of reduced coproducts. In this paper, the general formula is written explicitly in terms of Green functions and Feynman diagrams. The case we consider is that of a system with closed shells and a two-fold degenerate level occupied by a single electron. This does not look like a very interesting example, but we shall see that the hierarchy involves n -body Green functions with n up to 5. In the general case of a M -fold degenerate level, the Green functions involved go up to $n = 3M - 1$. In practice, such complicated Green functions are not usable, and the hierarchy is stopped after the first terms. The hierarchy we give here contains the first terms of the general hierarchy.

Quantum group concepts are not (yet) familiar in solid-state physics, so the pace will be slow and the calculations will be presented in full detail.

II. CALCULATION OF W_ρ^1

The simplest case is one electron ($N = 1$) in a twofold degenerate level ($M = 2$). The two degenerate levels are denoted by 1 and 2. The density matrix is

$$\rho = \begin{pmatrix} \rho_{11} & \rho_{12} \\ \rho_{21} & \rho_{22} \end{pmatrix},$$

with $\text{tr}(\rho) = \rho_{11} + \rho_{22} = 1$.

The moment generating function $\rho(\bar{\alpha}, \alpha)$ is defined in equation (28) of I. In our case, $N = 1$ and $\rho(\bar{\alpha}, \alpha) = \rho_0(\bar{\alpha}, \alpha) + \rho_1(\bar{\alpha}, \alpha)$. According to equation (29) of I (with $N = 1$, $i_1 = 1$ or 2, $j_1 = 1$ or 2)

$$\rho_1(\bar{\alpha}, \alpha) = \rho_{11}\bar{\alpha}_1\alpha_1 + \rho_{12}\bar{\alpha}_1\alpha_2 + \rho_{21}\bar{\alpha}_2\alpha_1 + \rho_{22}\bar{\alpha}_2\alpha_2.$$

Using now equation (30) of I with $k = 1$ we obtain

$$\rho_0(\bar{\alpha}, \alpha) = \text{tr}(\rho) = \rho_{11} + \rho_{22} = 1.$$

According to equation (32) of I, the cumulant generating function $\rho^c(\bar{\alpha}, \alpha)$ is obtained from

$$\log(\rho(\bar{\alpha}, \alpha)) = \log(1 + \rho_1(\bar{\alpha}, \alpha)) = \rho_1(\bar{\alpha}, \alpha) + \rho^c(\bar{\alpha}, \alpha),$$

where

$$\rho^c(\bar{\alpha}, \alpha) = \rho_2^c(\bar{\alpha}, \alpha) = -\frac{\rho_1(\bar{\alpha}, \alpha)^2}{2} = -\det(\rho)\bar{\alpha}_1\alpha_1\bar{\alpha}_2\alpha_2 = -(\rho_{11}\rho_{22} - \rho_{12}\rho_{21})\bar{\alpha}_1\alpha_1\bar{\alpha}_2\alpha_2.$$

According to equation (34) of I, this enables us to calculate W_ρ^0 . But we saw in section 7.3 of I that $W_\rho^1 = W_\rho^0$, thus

$$W_\rho^1 = -i \int \bar{\eta}(x) G_\rho^0(x, y) \eta(y) dx dy + \rho_2^c(\bar{\alpha}, \alpha), \quad (1)$$

with

$$G_\rho^0(x, y) = G_0^0(x, y) + i \left(\sum_{m=1}^C u_m(x) \bar{u}_m(y) + \sum_{k=1}^2 \sum_{l=1}^2 \rho_{lk} u_k(x) \bar{u}_l(y) \right) M,$$

where

$$M = \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}.$$

The expression for the 2×2 matrix $G_0^0(x, y)$ is given in equation (15) of I.

For later convenience, we give the identities

$$\begin{aligned} \frac{\delta^2 \bar{\alpha}_1 \alpha_1 \bar{\alpha}_2 \alpha_2}{\delta \eta_{\epsilon'}(y) \delta \bar{\eta}_\epsilon(x)} &= \epsilon \epsilon' (u_1(x) \bar{\alpha}_2 - u_2(x) \bar{\alpha}_1) (\bar{u}_1(y) \alpha_2 - \bar{u}_2(y) \alpha_1). \\ \frac{\delta^4 \rho_2^c(\bar{\alpha}, \alpha)}{\delta \bar{\eta}_{\epsilon_1}(x_1) \delta \bar{\eta}_{\epsilon_2}(x_2) \delta \eta_{\epsilon'_1}(y_1) \delta \eta_{\epsilon'_2}(y_2)} &= -\epsilon_1 \epsilon_2 \epsilon'_1 \epsilon'_2 r(x_1, x_2, y_1, y_2), \end{aligned} \quad (2)$$

where

$$r(x_1, x_2, y_1, y_2) = -\det(\boldsymbol{\rho})(u_1(x_1)u_2(x_2) - u_2(x_1)u_1(x_2))(\bar{u}_1(y_1)\bar{u}_2(y_2) - \bar{u}_2(y_1)\bar{u}_1(y_2)).$$

III. HIERARCHY

For one electron in a two-fold degenerate state, equation (43) of I stops at $n = 3$ because $M = 2$ and we have

$$\begin{aligned} \frac{\delta Z_\rho}{\delta \beta} &= \frac{\delta W_\rho^1}{\delta \beta} Z_\rho - i \sum (-1)^{|D_{(1')}|} (D_{(1')} \frac{\delta W_\rho^1}{\delta \beta}) (D_{(2')} Z_\rho) - \frac{1}{2} \sum (-1)^{|D_{(1')}^2|} (D_{(1')}^2 \frac{\delta W_\rho^1}{\delta \beta}) (D_{(2')}^2 Z_\rho) \\ &\quad + \frac{i}{6} \sum (-1)^{|D_{(1')}^3|} (D_{(1')}^3 \frac{\delta W_\rho^1}{\delta \beta}) (D_{(2')}^3 Z_\rho). \end{aligned}$$

To obtain the Green function, we put $\beta = \bar{\eta}_\epsilon(x)$, we make a functional derivative with respect to $\eta_{\epsilon'}(y)$ and we set all fermion sources to zero (in particular, this implies $Z_\rho = 1$). We use also the fact that a functional derivative of Z_ρ or W_ρ^1 with respect to an odd number of fermion sources is zero if the sources are set to zero. This enables us to calculate the signs $(-1)^{|D_{(1')}^n|}$.

$$\begin{aligned} \frac{\delta^2 Z_\rho}{\delta \eta_{\epsilon'}(y) \delta \bar{\eta}_\epsilon(x)} &= \frac{\delta^2 W_\rho^1}{\delta \eta_{\epsilon'}(y) \delta \bar{\eta}_\epsilon(x)} - i \sum (D_{(1')} \frac{\delta^2 W_\rho^1}{\delta \eta_{\epsilon'}(y) \delta \bar{\eta}_\epsilon(x)}) (D_{(2')} Z_\rho) - i \sum (D_{(1')} \frac{\delta W_\rho^1}{\delta \bar{\eta}_\epsilon(x)}) (D_{(2')} \frac{\delta Z_\rho}{\delta \eta_{\epsilon'}(y)}) \\ &\quad - \frac{1}{2} \sum (D_{(1')}^2 \frac{\delta^2 W_\rho^1}{\delta \eta_{\epsilon'}(y) \delta \bar{\eta}_\epsilon(x)}) (D_{(2')}^2 Z_\rho) - \frac{1}{2} \sum (D_{(1')}^2 \frac{\delta W_\rho^1}{\delta \bar{\eta}_\epsilon(x)}) (D_{(2')}^2 \frac{\delta Z_\rho}{\delta \eta_{\epsilon'}(y)}) \\ &\quad + \frac{i}{6} \sum (D_{(1')}^3 \frac{\delta W_\rho^1}{\delta \bar{\eta}_\epsilon(x)}) (D_{(2')}^3 \frac{\delta Z_\rho}{\delta \eta_{\epsilon'}(y)}). \end{aligned} \quad (3)$$

A. Coproduct

The non-relativistic interacting Hamiltonian is

$$\int H^{\text{int}}(t) dt = \frac{e^2}{2} \int \bar{\psi}(t, \mathbf{r}) \bar{\psi}(t, \mathbf{r}') \frac{1}{|\mathbf{r} - \mathbf{r}'|} \psi(t, \mathbf{r}') \psi(t, \mathbf{r}) dt d\mathbf{r} d\mathbf{r}'. \quad (4)$$

To simplify the notation, we define $x_1 = (t, \mathbf{r})$, $x'_1 = (t', \mathbf{r}')$ and $v(x_1 - x'_1) = \delta(t - t') e^2 / |\mathbf{r} - \mathbf{r}'|$, so that

$$\int H^{\text{int}}(t) dt = \frac{1}{2} \int v(x_1 - x'_1) \bar{\psi}(x_1) \bar{\psi}(x'_1) \psi(x'_1) \psi(x_1) dx_1 dx'_1.$$

To define the interacting operator D (i.e. the functional differential form of $\int H^{\text{int}}(t) dt$), we make the substitutions $\psi(x_1) \rightarrow -i\delta/\delta\bar{\eta}(x_1)$ and $\bar{\psi}(x_1) \rightarrow i\delta/\delta\eta(x_1)$ and we use the simplified notation

$$\delta_{1\pm} = \frac{\delta}{\delta\bar{\eta}_\pm(x_1)}, \quad \delta_{\bar{1}\pm} = \frac{\delta}{\delta\eta_\pm(x_1)}, \quad \delta_{1'\pm} = \frac{\delta}{\delta\bar{\eta}_\pm(x'_1)}, \quad \delta_{\bar{1}'\pm} = \frac{\delta}{\delta\eta_\pm(x'_1)}.$$

This gives us $D = D_+ - D_-$ with

$$D_{\pm} = \frac{1}{2} \int v(x_1 - x'_1) \delta_{1\pm} \delta_{1'\pm} \delta_{\bar{1}\pm} \delta_{\bar{1}'\pm} dx_1 dx'_1. \quad (5)$$

Thus

$$D = \frac{1}{2} \int dx_1 dx'_1 v(x_1 - x'_1) (\delta_{1+} \delta_{1'+} \delta_{\bar{1}+} \delta_{\bar{1}'+} - \delta_{1-} \delta_{1'-} \delta_{\bar{1}-} \delta_{\bar{1}'-}),$$

To further simplify the notation, we replace $\delta_{1\pm}$ by δ_1 and x_1 by 1 and x'_1 by $1'$. The interacting operator becomes

$$D = \frac{1}{2} \int d1 d1' v(1 - 1') \sum_{\pm} \pm \delta_1 \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}}.$$

To calculate the coproduct of D , we start from the coproduct of the basic functional derivatives

$$\Delta \delta_1 = \delta_1 \otimes \mathbf{1} + \mathbf{1} \otimes \delta_1, \quad \Delta \delta_{1'} = \delta_{1'} \otimes \mathbf{1} + \mathbf{1} \otimes \delta_{1'}, \quad \Delta \delta_{\bar{1}} = \delta_{\bar{1}} \otimes \mathbf{1} + \mathbf{1} \otimes \delta_{\bar{1}}, \quad \Delta \delta_{\bar{1}'} = \delta_{\bar{1}'} \otimes \mathbf{1} + \mathbf{1} \otimes \delta_{\bar{1}}'.$$

Then we use equation (21) of I to calculate the coproduct of $\delta_1 \delta_{1'}$ and $\delta_{\bar{1}} \delta_{\bar{1}'}$:

$$\Delta \delta_1 \delta_{1'} = \delta_1 \delta_{1'} \otimes \mathbf{1} + \delta_1 \otimes \delta_{1'} - \delta_{1'} \otimes \delta_1 + \mathbf{1} \otimes \delta_1 \delta_{1'}, \quad \Delta \delta_{\bar{1}} \delta_{\bar{1}'} = \delta_{\bar{1}} \delta_{\bar{1}'} \otimes \mathbf{1} + \delta_{\bar{1}'} \otimes \delta_{\bar{1}} - \delta_{\bar{1}} \otimes \delta_{\bar{1}'} + \mathbf{1} \otimes \delta_{\bar{1}} \delta_{\bar{1}'}.$$

We use equation (21) of I again to obtain

$$\begin{aligned} \Delta \delta_1 \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}'} &= \delta_1 \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}'} \otimes \mathbf{1} + \mathbf{1} \otimes \delta_1 \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}'} + \delta_1 \delta_{1'} \delta_{\bar{1}} \otimes \delta_{\bar{1}'} - \delta_1 \delta_{1'} \delta_{\bar{1}} \otimes \delta_{\bar{1}'} + \delta_1 \delta_{1'} \otimes \delta_{\bar{1}} \delta_{\bar{1}'} + \delta_1 \delta_{\bar{1}} \delta_{\bar{1}'} \otimes \delta_{1'} \\ &\quad - \delta_1 \delta_{\bar{1}'} \otimes \delta_{1'} \delta_{\bar{1}} + \delta_1 \delta_{\bar{1}} \otimes \delta_{1'} \delta_{\bar{1}'} + \delta_1 \otimes \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}'} - \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}'} \otimes \delta_1 + \delta_{1'} \delta_{\bar{1}'} \otimes \delta_1 \delta_{\bar{1}} - \delta_{1'} \delta_{\bar{1}} \otimes \delta_1 \delta_{\bar{1}'} \\ &\quad - \delta_{1'} \otimes \delta_1 \delta_{\bar{1}} \delta_{\bar{1}'} + \delta_{\bar{1}} \delta_{\bar{1}'} \otimes \delta_1 \delta_{1'} + \delta_{\bar{1}'} \otimes \delta_1 \delta_{1'} \delta_{\bar{1}} - \delta_{\bar{1}} \otimes \delta_1 \delta_{1'} \delta_{\bar{1}'} \end{aligned} \quad (6)$$

The reduced coproduct with respect to D is obtained by its definition $\Delta' D = \Delta D - D \otimes \mathbf{1} - \mathbf{1} \otimes D$. Therefore,

$$\begin{aligned} \Delta' D &= \frac{1}{2} \int d1 d1' v(1 - 1') \sum_{\pm} \pm \left(\delta_1 \delta_{1'} \delta_{\bar{1}} \otimes \delta_{\bar{1}'} - \delta_1 \delta_{1'} \delta_{\bar{1}} \otimes \delta_{\bar{1}'} + \delta_1 \delta_{1'} \otimes \delta_{\bar{1}} \delta_{\bar{1}'} + \delta_1 \delta_{\bar{1}} \delta_{\bar{1}'} \otimes \delta_{1'} \right. \\ &\quad \left. - \delta_1 \delta_{\bar{1}'} \otimes \delta_{1'} \delta_{\bar{1}} + \delta_1 \delta_{\bar{1}} \otimes \delta_{1'} \delta_{\bar{1}'} + \delta_1 \otimes \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}'} - \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}'} \otimes \delta_1 + \delta_{1'} \delta_{\bar{1}'} \otimes \delta_1 \delta_{\bar{1}} - \delta_{1'} \delta_{\bar{1}} \otimes \delta_1 \delta_{\bar{1}'} \right. \\ &\quad \left. - \delta_{1'} \otimes \delta_1 \delta_{\bar{1}} \delta_{\bar{1}'} + \delta_{\bar{1}} \delta_{\bar{1}'} \otimes \delta_1 \delta_{1'} + \delta_{\bar{1}'} \otimes \delta_1 \delta_{1'} \delta_{\bar{1}} - \delta_{\bar{1}} \otimes \delta_1 \delta_{1'} \delta_{\bar{1}'} \right). \end{aligned} \quad (7)$$

Notice that, in $\Delta' D = \sum D_{(1')} \otimes D_{(2')}$, each term $D_{(1')}$ or $D_{(2')}$ contains between one and three functional derivatives.

B. Expansion of equation (3)

Now we are going to examine each term of equation (3). We define the n -body Green function as

$$G_{\epsilon_1 \dots \epsilon_n \epsilon'_1 \dots \epsilon'_n}(x_1, \dots, x_n, y'_1, \dots, y'_n) = (-i)^n \frac{\delta^{2n} Z_{\rho}}{\delta \bar{\eta}_{\epsilon_1}(x_1) \dots \delta \bar{\eta}_{\epsilon_n}(x_n) \delta \eta_{\epsilon'_1}(y_1) \dots \delta \eta_{\epsilon'_n}(y_n)}. \quad (8)$$

The right hand side is taken at zero external sources. To simplify the notation, we write i for x_i and i' for y_i , we consider that ϵ_i goes with the variable x_i , ϵ'_i with y_i , so the ϵ are now implicit and we have

$$G(1, \dots, n, 1', \dots, n') = (-i)^n \delta_{\bar{1}} \dots \delta_{\bar{n}} \delta_{1'} \dots \delta_{n'} Z_{\rho}.$$

With this notation, the left hand side of equation (3) is now

$$\frac{\delta^2 Z_{\rho}}{\delta \eta_{\epsilon'}(y) \delta \bar{\eta}_{\epsilon}(x)} = - \frac{\delta^2 Z_{\rho}}{\delta \bar{\eta}_{\epsilon}(x) \delta \eta_{\epsilon'}(y)} = -i G(x, y).$$

More explicitly $G(x, y) = G_{\epsilon \epsilon'}(x, y)$. In the following we shall calculate in detail all the terms of the right hand side of equation (3).

C. First term of equation (3)

The first term of (3) is very simple. From equation (1), we know that W_ρ^1 is the sum of a term of degree two and a term of degree four in the external sources. The first term in the right hand side of equation (3) contains only two functional derivatives. Thus, at zero sources, the second term of W_ρ^1 does not contribute. By definition of functional derivatives, the first term gives us

$$\frac{\delta^2 W_\rho^1}{\delta \eta_{\epsilon'}(y) \delta \bar{\eta}_\epsilon(x)} = -i G_{\rho \epsilon \epsilon'}^0(x, y).$$

D. Second term of equation (3)

The second term of equation (3) is

$$-i \sum (D_{(1')} \delta_y \delta_{\bar{x}} W_\rho^1) (D_{(2')} Z_\rho),$$

where we wrote δ_y for $\delta/\delta \eta_{\epsilon'}(y)$ and $\delta_{\bar{x}}$ for $\delta/\delta \bar{\eta}_\epsilon(x)$. According to equations (7), each term $D_{(1')}$ contains at least one functional derivative. Therefore, the first term of W_ρ^1 does not contribute because it is of degree 2. The second term of W_ρ^1 is of degree four, and it contains two sources η and two sources $\bar{\eta}$. The functional derivatives are evaluated at zero sources, thus $D_{(1')}$ must be the product of a derivative with respect to a source η and a derivative with respect to a source $\bar{\eta}$. According to equation (7), there are four terms with this property in $\Delta' D$:

$$\frac{1}{2} \int d1 d1' v(1-1') \sum_{\pm} \pm \left(-\delta_1 \delta_{\bar{1}'} \otimes \delta_{1'} \delta_{\bar{1}} + \delta_1 \delta_{\bar{1}} \otimes \delta_{1'} \delta_{\bar{1}'} + \delta_{1'} \delta_{\bar{1}'} \otimes \delta_1 \delta_{\bar{1}} - \delta_{1'} \delta_{\bar{1}} \otimes \delta_1 \delta_{\bar{1}'} \right).$$

The second term of equation (3) becomes

$$\begin{aligned} -\frac{i}{2} \int d1 d1' v(1-1') \sum_{\pm} \pm \left(-(\delta_1 \delta_{\bar{1}'} \delta_y \delta_{\bar{x}} W_\rho^1) (\delta_{1'} \delta_{\bar{1}} Z_\rho) + (\delta_1 \delta_{\bar{1}} \delta_y \delta_{\bar{x}} W_\rho^1) (\delta_{1'} \delta_{\bar{1}'} Z_\rho) + (\delta_{1'} \delta_{\bar{1}'} \delta_y \delta_{\bar{x}} W_\rho^1) (\delta_1 \delta_{\bar{1}} Z_\rho) \right. \\ \left. - (\delta_{1'} \delta_{\bar{1}} \delta_y \delta_{\bar{x}} W_\rho^1) (\delta_1 \delta_{\bar{1}'} Z_\rho) \right). \end{aligned} \quad (9)$$

Let us now consider the first term of (9). According to (8), $\delta_{1'} \delta_{\bar{1}} Z_\rho = -i G(1, 1')$. According to (2),

$$\delta_1 \delta_{\bar{1}'} \delta_y \delta_{\bar{x}} W_\rho^1 = \delta_{\bar{x}} \delta_{\bar{1}'} \delta_1 \delta_y W_\rho^1 = -\epsilon \epsilon' r(x, 1', 1, y).$$

For the second term of (9) we have $\delta_{1'} \delta_{\bar{1}'} Z_\rho = -i G(1', 1')$ and

$$\delta_1 \delta_{\bar{1}} \delta_y \delta_{\bar{x}} W_\rho^1 = \delta_{\bar{x}} \delta_{\bar{1}} \delta_1 \delta_y W_\rho^1 = -\epsilon \epsilon' r(x, 1, 1, y).$$

Interchanging the variables 1 and $1'$, we see that the third term of (9) is equal to the first one, and the fourth term is equal to the second one. Finally, the second term of (3) becomes the sum of

$$-\epsilon \epsilon' \int d1 d1' v(1-1') r(x, 1', 1, y) \sum_{\pm} \pm G(1, 1'),$$

and

$$\epsilon \epsilon' \int d1 d1' v(1-1') r(x, 1, 1, y) \sum_{\pm} \pm G(1', 1').$$

E. Third term of equation (3)

The third term is

$$-i \sum (D_{(1')} \delta_{\bar{x}} W_\rho^1) (D_{(2')} \delta_y Z_\rho).$$

For this term, two kinds of $D_{(1')} \otimes D_{(2')}$ intervene: with one or three products in $D_{(1')}$. For the first kind of terms, the relevant coproducts are $\delta_1 \otimes \delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}} - \delta_{1'} \otimes \delta_1 \delta_{\bar{1}} \delta_{\bar{1}}$ and the result is

$$\begin{aligned} & -\frac{i}{2} \int d1 d1' v(1-1') \sum_{\pm} \pm \left((\delta_1 \delta_{\bar{x}} W_{\rho}^1) (\delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}} \delta_y Z_{\rho}) - (\delta_{1'} \delta_{\bar{x}} W_{\rho}^1) (\delta_1 \delta_{\bar{1}} \delta_{\bar{1}} \delta_y Z_{\rho}) \right) \\ & = -i \int d1 d1' v(1-1') \sum_{\pm} \pm (\delta_1 \delta_{\bar{x}} W_{\rho}^1) (\delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}} \delta_y Z_{\rho}). \end{aligned}$$

We have interchanged the variables 1 and 1' to obtain the last line. We use now $\delta_1 \delta_{\bar{x}} W_{\rho}^1 = -i G_{\rho}^0(x, 1)$ and

$$\delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}} \delta_y Z_{\rho} = -\delta_{\bar{1}} \delta_{1'} \delta_{\bar{1}} \delta_y Z_{\rho} = G(1, 1', 1', y),$$

to get

$$- \int d1 d1' v(1-1') \sum_{\pm} \pm G_{\rho}^0(x, 1) G(1, 1', 1', y).$$

For the second kind of terms, the relevant coproducts are $\delta_1 \delta_{1'} \delta_{\bar{1}} \otimes \delta_{\bar{1}} - \delta_1 \delta_{1'} \delta_{\bar{1}} \otimes \delta_{\bar{1}}$. Again, interchanging 1 and 1' shows that these two terms give the same contribution and the result is

$$i \int d1 d1' v(1-1') \sum_{\pm} \pm (\delta_1 \delta_{1'} \delta_{\bar{1}} \delta_{\bar{x}} W_{\rho}^1) (\delta_{\bar{1}} \delta_y Z_{\rho}).$$

We calculate

$$\delta_1 \delta_{1'} \delta_{\bar{1}} \delta_{\bar{x}} W_{\rho}^1 = -\delta_{\bar{x}} \delta_{\bar{1}} \delta_1 \delta_{1'} W_{\rho}^1 = \pm \epsilon r(x, 1, 1, 1'),$$

and $\delta_{\bar{1}} \delta_y Z_{\rho} = i G(1', y)$. The contribution of the second kind of terms becomes

$$-\epsilon \int d1 d1' v(1-1') r(x, 1, 1, 1') \sum_{\pm} G(1', y).$$

F. Fourth term of equation (3)

The fourth term of equation (3) is

$$-\frac{1}{2} \sum (D_{(1')}^2 \delta_y \delta_{\bar{x}} W_{\rho}^1) (D_{(2')}^2 Z_{\rho}).$$

Here we meet something new because we must use the reduced coproduct $\Delta' D^2$. According to the general formula (39) of I, this reduced coproduct is obtained by multiplying $\Delta' D$ by itself. The spacetime variables of the first $\Delta' D$ will still be denoted by 1 and 1', while the spacetime variables of the second $\Delta' D$ will be denoted by 2 and 2'. Moreover, the $\pm$ variable of the second $\Delta' D$ will be written $\pm'$. Each term $D_{(1')}$ of $\Delta' D$ is at least of degree 1, so each term of $D_{(1')}^2$ is at least of degree 2. Therefore, the only term of W_{ρ}^1 that contributes to $D_{(1')}^2 \delta_y \delta_{\bar{x}} W_{\rho}^1$ is $\rho^c(\bar{\alpha}, \alpha)$. We use again the fact that $\rho^c(\bar{\alpha}, \alpha)$ is a product of two sources η and two sources $\bar{\eta}$ to deduce that $D_{(1')}^2$ must be the product of a functional derivative with respect to η and a functional derivative with respect to $\bar{\eta}$. The eight terms $D_{(1')}^2$ that satisfy this condition are $\delta_1 \delta_{\bar{2}}$, $\delta_1 \delta_{\bar{2}'}$, $\delta_{1'} \delta_{\bar{2}}$, $\delta_{1'} \delta_{\bar{2}'}$, $\delta_{\bar{1}} \delta_2$, $\delta_{\bar{1}} \delta_{2'}$, $\delta_{\bar{1}} \delta_{2'}$, $\delta_{\bar{1}} \delta_2$ and $\delta_{\bar{1}} \delta_{2'}$. This gives us eight terms which are all identical if we interchange 1 with 1' or 2 with 2' or (1,1') with (2,2'). Thus, the fourth term of equation (3) becomes

$$- \int d1 d1' d2 d2' v(1-1') v(2-2') \sum_{\pm \pm'} \pm \pm' (\delta_1 \delta_{\bar{2}} \delta_y \delta_{\bar{x}} W_{\rho}^1) (\delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}} \delta_2 \delta_{2'} \delta_{\bar{2}'} Z_{\rho}).$$

We have just to calculate

$$\delta_1 \delta_{\bar{2}} \delta_y \delta_{\bar{x}} W_{\rho}^1 = \delta_{\bar{x}} \delta_{\bar{2}} \delta_1 \delta_y W_{\rho}^1 = -\epsilon \epsilon' \pm \pm' r(x, 2, 1, y),$$

and

$$\delta_{1'} \delta_{\bar{1}} \delta_1 \delta_2 \delta_{2'} \delta_{\bar{2}'} Z_{\rho} = \delta_{\bar{2}'} \delta_{\bar{1}} \delta_{1'} \delta_2 \delta_{2'} \delta_{1'} Z_{\rho} = -i G(2', 1, 1', 2, 2', 1').$$

The final expression for the fourth term is then

$$-i \epsilon \epsilon' \int d1 d1' d2 d2' v(1-1') v(2-2') r(x, 2, 1, y) \sum_{\pm \pm'} G(2', 1, 1', 2, 2', 1').$$

G. Fifth term of equation (3)

The fifth term of equation (3) is

$$-\frac{1}{2} \sum (D_{(1')}^2 \delta_{\bar{x}} W_\rho^1) (D_{(2')}^2 \delta_y Z_\rho).$$

To give nonzero contributions, $D_{(1')}^2$ must contain three functional derivatives: one functional derivative with respect to a $\bar{\eta}$ and two with respect to η . This gives us three kinds of terms: (i) the argument of $\bar{\eta}$ does not come from the same D as the arguments of the two η , as in $D_{(1')}^2 = \delta_{\bar{1}} \delta_2 \delta_{2'}$ (ii) the argument of $\bar{\eta}$ is the same as that of one of the two η , for example $D_{(1')}^2 = \delta_1 \delta_{2'} \delta_{\bar{2}'}$ (iii) the argument of $\bar{\eta}$ comes from the same D as the argument of one of the two η but they are not identical, for example $D_{(1')}^2 = -\delta_1 \delta_2 \delta_{\bar{2}'}$. In $\Delta' D^2$ there are four terms of type (i), eight terms of type (ii) and eight terms of type (iii). All the terms of each class give the same contribution. We calculate now the contribution of the three kinds of terms

1. Terms of kind (i)

The sum of the terms of type (i) is

$$-\frac{1}{2} \int d1 d1' d2 d2' v(1-1') v(2-2') \sum_{\pm, \pm'} \pm \pm' (\delta_{\bar{1}} \delta_2 \delta_{2'} \delta_{\bar{x}} W_\rho^1) (\delta_1 \delta_{1'} \delta_{\bar{1}} \delta_{\bar{2}'} \delta_y Z_\rho).$$

We compute

$$\delta_{\bar{1}} \delta_2 \delta_{2'} \delta_{\bar{x}} W_\rho^1 = -\delta_{\bar{x}} \delta_{\bar{1}} \delta_2 \delta_{2'} W_\rho^1 = \pm \epsilon r(x, 1', 2, 2'),$$

and

$$\delta_1 \delta_{1'} \delta_{\bar{1}} \delta_{\bar{2}'} \delta_y Z_\rho = -\delta_{\bar{1}} \delta_2 \delta_{2'} \delta_1 \delta_{1'} \delta_y Z_\rho = iG(1, 2, 2', 1, 1', y)$$

to obtain

$$-\frac{i}{2} \epsilon \int d1 d1' d2 d2' v(1-1') v(2-2') r(x, 1', 2, 2') \sum_{\pm, \pm'} \pm' G(1, 2, 2', 1, 1', y).$$

2. Terms of kind (ii)

The sum of the terms of type (ii) is

$$-\int d1 d1' d2 d2' v(1-1') v(2-2') \sum_{\pm, \pm'} \pm \pm' (\delta_1 \delta_{2'} \delta_{\bar{2}'} \delta_{\bar{x}} W_\rho^1) (\delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}} \delta_2 \delta_{\bar{2}'} \delta_y Z_\rho).$$

We use

$$\delta_1 \delta_{2'} \delta_{\bar{2}'} \delta_{\bar{x}} W_\rho^1 = -\delta_{\bar{x}} \delta_{\bar{2}'} \delta_1 \delta_{2'} W_\rho^1 = \pm \epsilon r(x, 2', 1, 2'),$$

and

$$\delta_{1'} \delta_{\bar{1}} \delta_{\bar{1}} \delta_2 \delta_{\bar{2}'} \delta_y Z_\rho = -\delta_{\bar{1}} \delta_{\bar{1}} \delta_{\bar{2}'} \delta_1 \delta_{2'} \delta_y Z_\rho = iG(1, 1', 2, 1', 2, y).$$

Therefore, the fifth term of type (ii) becomes

$$-i\epsilon \int d1 d1' d2 d2' v(1-1') v(2-2') r(x, 2', 1, 2') \sum_{\pm, \pm'} \pm' G(1, 1', 2, 1', 2, y).$$

3. Terms of kind (iii)

The sum of the terms of type (iii) is

$$\int d1d1'd2d2'v(1-1')v(2-2') \sum_{\pm, \pm'} \pm \pm' (\delta_1 \delta_2 \delta_{\bar{2}'} \delta_{\bar{x}} W_\rho^1) (\delta_{1'} \delta_{\bar{1}'} \delta_{\bar{1}} \delta_{2'} \delta_{\bar{2}} \delta_y Z_\rho).$$

We have

$$\delta_1 \delta_2 \delta_{\bar{2}'} \delta_{\bar{x}} W_\rho^1 = \delta_{\bar{x}} \delta_{\bar{2}'} \delta_1 \delta_2 W_\rho^1 = \mp \epsilon r(x, 2', 1, 2),$$

and

$$\delta_{1'} \delta_{\bar{1}'} \delta_{\bar{1}} \delta_{2'} \delta_{\bar{2}} \delta_y Z_\rho = -\delta_{\bar{1}} \delta_{\bar{1}'} \delta_{\bar{2}} \delta_{1'} \delta_{2'} \delta_y Z_\rho = iG(1, 1', 2, 1', 2', y).$$

The fifth term of type (iii) becomes

$$-i\epsilon \int d1d1'd2d2'v(1-1')v(2-2')r(x, 2', 1, 2) \sum_{\pm, \pm'} \pm' G(1, 1', 2, 1', 2', y).$$

H. Sixth term of equation (3)

The sixth term of equation (3) is

$$\frac{i}{6} \sum (D_{(1')}^3 \delta_{\bar{x}} W_\rho^1) (D_{(2')}^3 \delta_y Z_\rho).$$

It looks complicated because we have now $\Delta' D^3 = (\Delta' D)(\Delta' D)(\Delta' D)$. The third $\Delta' D$ will have the variables 3, 3' and $\pm''$. Again, we need one derivative with respect to $\bar{\eta}$ and two derivatives with respect to η in $D_{(1')}^3$. There are 24 terms of this kind in $\Delta' D^3$, and they all give the same contribution as $\delta_{\bar{1}} \delta_2 \delta_3 \otimes \delta_1 \delta_{1'} \delta_{\bar{1}'} \delta_{2'} \delta_{\bar{2}'} \delta_{\bar{2}} \delta_{3'} \delta_{\bar{3}'} \delta_{\bar{3}}$. Thus, the sixth term is

$$\frac{i}{2} \int d1d1'd2d2'd3d3'v(1-1')v(2-2')v(3-3') \sum_{\pm \pm' \pm''} \pm \pm' \pm'' (\delta_{\bar{1}} \delta_2 \delta_3 \delta_{\bar{x}} W_\rho^1) (\delta_{1'} \delta_{\bar{1}'} \delta_{2'} \delta_{\bar{2}'} \delta_{\bar{2}} \delta_{3'} \delta_{\bar{3}'} \delta_{\bar{3}} \delta_y Z_\rho).$$

We compute

$$\delta_{\bar{1}} \delta_2 \delta_3 \delta_{\bar{x}} W_\rho^1 = -\delta_{\bar{x}} \delta_{\bar{1}} \delta_2 \delta_3 W_\rho^1 = \pm \pm' \pm'' \epsilon r(x, 1, 2, 3),$$

and

$$\delta_1 \delta_{1'} \delta_{\bar{1}'} \delta_{2'} \delta_{\bar{2}'} \delta_{\bar{2}} \delta_{3'} \delta_{\bar{3}'} \delta_{\bar{3}} \delta_y Z_\rho = \delta_{\bar{1}} \delta_{\bar{2}} \delta_{2'} \delta_{\bar{3}} \delta_{\bar{3}'} \delta_1 \delta_{1'} \delta_{2'} \delta_{3'} \delta_y Z_\rho = iG(1', 2, 2', 3, 3', 1, 1', 2', 3', y).$$

The last term of equation (3) is then

$$-\frac{1}{2} \epsilon \int d1d1'd2d2'd3d3'v(1-1')v(2-2')v(3-3')r(x, 1, 2, 3) \sum_{\pm \pm' \pm''} G(1', 2, 2', 3, 3', 1, 1', 2', 3', y).$$




FIG. 1: Diagrammatic representation of equation (10).

I. The Green function hierarchy

If we gather all the previous results, we obtain the Green function hierarchy

$$\begin{aligned}
G(x, y) &= G_{\epsilon\epsilon'}(x, y) = G_{\rho\epsilon\epsilon'}^0(x, y) - i \int d1d1'v(1-1') \sum_{\pm} \pm G_{\rho}^0(x, 1)G(1, 1', 1', y) \\
&\quad - i\epsilon\epsilon' \int d1d1'v(1-1')r(x, 1', 1, y) \sum_{\pm} \pm G(1, 1') + i\epsilon\epsilon' \int d1d1'v(1-1')r(x, 1, 1, y) \sum_{\pm} \pm G(1', 1') \\
&\quad - i\epsilon \int d1d1'v(1-1')r(x, 1, 1, 1') \sum_{\pm} G(1', y) \\
&\quad + \frac{1}{2}\epsilon \int d1d1'd2d2'v(1-1')v(2-2')r(x, 1', 2, 2') \sum_{\pm, \pm'} \pm' G(1, 2, 2', 1, 1', y) \\
&\quad + \epsilon \int d1d1'd2d2'v(1-1')v(2-2')r(x, 2', 1, 2) \sum_{\pm, \pm'} \pm' G(1, 1', 2, 1', 2', y). \\
&\quad + \epsilon\epsilon' \int d1d1'd2d2'v(1-1')v(2-2')r(x, 2, 1, y) \sum_{\pm\pm'} G(2', 1, 1', 2, 2', 1') \\
&\quad + \epsilon \int d1d1'd2d2'v(1-1')v(2-2')r(x, 2', 1, 2') \sum_{\pm, \pm'} \pm' G(1, 1', 2, 1', 2, y). \\
&\quad - \frac{i}{2}\epsilon \int d1d1'd2d2'd3d3'v(1-1')v(2-2')v(3-3')r(x, 1, 2, 3) \sum_{\pm\pm'\pm''} G(1', 2, 2', 3, 3', 1, 1', 2', 3', y). \quad (10)
\end{aligned}$$

In terms of the Feynman diagrams of nonequilibrium quantum field theory², these terms can be rewritten as in figure 1. The order of the terms is the same in equation (10) and in figure 1.

IV. CONCLUSION

The hierarchy of Green functions presented here is an important step in the effective calculation of degenerate systems. However, the practical implementation of this hierarchy requires an approximation to close it. The GW approximation³⁻⁵ is a well-known and powerful way of closing the hierarchy. It has to be adapted to our more complicated hierarchy. For the calculation of the optical properties of allochromatic crystals, it will also be necessary to adapt the Bethe-Salpeter approach⁶ to our setting.

Moreover, a functional derivation of the energy with respect to the density matrix provides equations that enable us to unify the Green-function formalism and the diagonalization method of many-body theory. This will be presented in a forthcoming publication⁷.

References

- ¹ Ch. Brouder. Many-body theory of degenerate systems. 2003. arXiv:cond-mat/0309558.
- ² A.G. Hall. Non-equilibrium Green functions: Generalized Wick's theorem and diagrammatic perturbation theory with initial correlations. *J. Phys. A: Math. Gen.*, 8:214–25, 1975.
- ³ L. Hedin. New method for calculating the one-particle Green's function with application to the electron-gas problem. *Phys. Rev. A*, 139:796–823, 1965.
- ⁴ F. Aryasetiawan and O. Gunnarson. The GW method. *Rep. Prog. Phys.*, 61:237–312, 1998.
- ⁵ F. Aryasetiawan. The GW approximation and vertex corrections. In V.I. Anisimov, editor, *Strong Coulomb correlations in electronic structure calculations*, pages 1–95, Amsterdam, 2000. Gordon and Breach.
- ⁶ G. Onida, L. Reining, and A. Rubio. Electronic excitations: density-functional versus many-body Green's-function approaches. *Rev. Mod. Phys.*, 74:601–59, 2002.
- ⁷ Ch. Brouder. Many-body approach to crystal field theory. 2003. arXiv:cond-mat/0308058.